

SOME PROPERTIES OF GENERALIZED LOCAL COHOMOLOGY MODULES

AMIR MAFI

ABSTRACT. Let R be a commutative Noetherian ring, $\mathfrak{a}$ an ideal of R , M and N be two finitely generated R -modules. Let t be a positive integer. We prove that if R is local with maximal ideal $\mathfrak{m}$ and $M \otimes_R N$ is of finite length then $H_{\mathfrak{m}}^t(M, N)$ is of finite length for all $t \geq 0$ and $l_R(H_{\mathfrak{m}}^t(M, N)) \leq \sum_{i=0}^t l_R(\text{Ext}_R^i(M, H_{\mathfrak{m}}^{t-i}(N)))$. This yields, $l_R(H_{\mathfrak{m}}^t(M, N)) = l_R(\text{Ext}_R^t(M, N))$. Additionally, we show that $\text{Ext}_R^i(R/\mathfrak{a}, N)$ is Artinian for all $i \leq t$ if and only if $H_{\mathfrak{a}}^i(M, N)$ is Artinian for all $i \leq t$. Moreover, we show that whenever $\dim(R/\mathfrak{a}) = 0$ then $H_{\mathfrak{a}}^t(M, N)$ is Artinian for all $t \geq 0$.

1. INTRODUCTION

Throughout this paper, R is a commutative Noetherian ring with identity and $\mathfrak{a}$ is an ideal of R . For an R -module N and non-negative integer t , the local cohomology module $H_{\mathfrak{a}}^t(N)$ was first introduced and studied by Grothendieck [4]. For example he showed that $H_{\mathfrak{m}}^t(N)$ is Artinian for all t , whenever N is finitely generated and R is local with maximal ideal $\mathfrak{m}$. One of the general problem in local cohomology is to find when the local cohomology module $H_{\mathfrak{a}}^t(N)$ is Artinian (see [6, Problem 3]). In [7], Melkersson proved that for an R -module N , $H_{\mathfrak{a}}^i(N)$ is Artinian for all $i \leq t$ if and only if $\text{Ext}_R^i(R/\mathfrak{a}, N)$ is Artinian for all $i \leq t$. The latter result leads us to consider the same result for generalized local cohomology modules which was introduced and studied by Herzog [5] (see also [10]). For each $i \geq 0$, the generalized local cohomology functor $H_{\mathfrak{a}}^i(., .)$ is defined by

$$H_{\mathfrak{a}}^i(M, N) = \varinjlim_n \text{Ext}_R^i(M/\mathfrak{a}^n M, N)$$

for all R -modules M and N . Clearly, this is a generalization of the usual local cohomology module. The reader is referred to articles [2], [1] and [3] for more results

2000 *Mathematics Subject Classification.* 13D45, 13E10.

Key words and phrases. Generalized local cohomology, spectral sequences, artinian modules.

on generalized local cohomology. Our main purpose in this paper is to show the following.

Theorem 1.1. *Let $\mathfrak{a}$ be an ideal of R , M a finitely generated R -module, and let t be a positive integer.*

(α) *If R is local with maximal ideal $\mathfrak{m}$ and N is a finitely generated R -module such that $M \otimes_R N$ is of finite length, then $l_R(H_{\mathfrak{m}}^t(M, N)) \leq \sum_{i=0}^t l_R(\text{Ext}_R^i(M, H_{\mathfrak{m}}^{t-i}(N)))$ for any $t \geq 0$.*

(β) *Let N be an R -module. Then the following statements are equivalent:*

(i) *$\text{Ext}_R^i(R/\mathfrak{a}, N)$ is an Artinian R -module for all $i \leq t$.*

(ii) *$H_{\mathfrak{a}}^i(M, N)$ is an Artinian R -module for all $i \leq t$.*

Clearly (α) extends the result [9, Theorem 3.2] and (β) is equivalent with the main result [7, Theorem 1.2].

2. THE RESULTS

We start this section with the following lemma.

Lemma 2.1. *Let R be a local ring with maximal ideal $\mathfrak{m}$, M and N be two finitely generated R -modules such that $M \otimes_R N$ is of finite length. Then $H_{\mathfrak{m}}^t(M, N)$ is of finite length for all $t \geq 0$.*

Proof. Consider the Grothendieck spectral sequence [8, Theorem 11.38]

$$E_2^{p,q} := \text{Ext}_R^p(M, H_{\mathfrak{m}}^q(N)) \Longrightarrow_p H_{\mathfrak{m}}^{p+q}(M, N).$$

for $t \geq 0$, we have a finite filtration

$$0 = \phi^{t+1}H^t \subseteq \phi^t H^t \subseteq \cdots \subseteq \phi^1 H^t \subseteq \phi^0 H^t = H_{\mathfrak{m}}^t(M, N),$$

such that $E_{\infty}^{i,t-i} = \phi^i H^t / \phi^{i+1} H^t$ for all $0 \leq i \leq t$. Now, since $E_{\infty}^{i,j}$ is a homomorphic image of $E_2^{i,j}$ for all $i, j \geq 0$, by [9, Lemma 3.1], $E_2^{i,j}$ is of finite length for all $i, j \geq 0$. It therefore follows that $E_{\infty}^{i,j}$ is of finite length for all $i, j \geq 0$ and that $\phi^{t-i} H^t$ is of finite length for all $0 \leq i \leq t$. Hence, using the exact sequence

$$0 \longrightarrow \phi^1 H^t \longrightarrow H_{\mathfrak{m}}^t(M, N) \longrightarrow E_{\infty}^{0,t} \longrightarrow 0 \quad (0 \leq i \leq t)$$

we get $H_{\mathfrak{m}}^t(M, N)$ is of finite length for all $t \geq 0$. $\square$

The following theorem extends [9, Theorem 3.2].

Theorem 2.2. *Let R be a local ring with maximal ideal $\mathfrak{m}$ and let M and N be two finitely generated R -modules such that $M \otimes_R N$ is of finite length. Then $l_R(H_{\mathfrak{m}}^t(M, N)) \leq \sum_{i=0}^t l_R(\text{Ext}_R^i(M, H_{\mathfrak{m}}^{t-i}(N)))$ for all $t \geq 0$. Consequently, $l_R(H_{\mathfrak{m}}^t(M, N)) = l_R(\text{Ext}_R^t(M, N))$.*

Proof. Let $2 \leq i \leq t$. With the notation of [8, §11] we consider the exact sequences

$$0 \longrightarrow \ker d_i^{i,t-i} \longrightarrow E_i^{i,t-i} \longrightarrow E_i^{2i,t-2i+1}$$

and note that $E_i^{i,t-i} = \ker d_{i-1}^{i,t-i} / \text{im } d_{i-1}^{1,t-2}$ and $E_i^{i,j} = 0$ for all $j < 0$. So, we have $\ker d_{t+2}^{i,t-i} \cong E_{t+2}^{i,t-i} \cong \dots \cong E_{\infty}^{i,t-i}$ for all $(0 \leq i \leq t)$. Now, using the exact sequences

$$0 \longrightarrow \phi^{i+1}H^t \longrightarrow \phi^iH^t \longrightarrow E_{\infty}^{i,t-i} \longrightarrow 0 \quad (0 \leq i \leq t)$$

and an argument similar to that used in 1.1 together with the facts $E_{\infty}^{i,t-i} \cong \ker d_{t+2}^{i,t-i} \subseteq \ker d_2^{i,t-i} \subseteq E_2^{i,t-i}$ for all $0 \leq i \leq t$, one can deduce that $l_R(H_{\mathfrak{m}}^t(M, N)) \leq \sum_{i=0}^t l_R(\text{Ext}_R^i(M, H_{\mathfrak{m}}^{t-i}(N)))$. Moreover, there is a spectral sequence

$$E_2^{i,j} = \text{Ext}_R^i(M, H_{\mathfrak{m}}^j(N)) \implies_i \text{Ext}_R^{i+j}(M, N).$$

Hence, we have a finite filtration

$$0 = \psi^{t+1}H^t \subseteq \psi^tH^t \subseteq \dots \subseteq \psi^1H^t \subseteq \psi^0H^t = \text{Ext}_R^t(M, N),$$

such that $E_{\infty}^{i,t-i} = \psi^iH^t / \psi^{i+1}H^t$ for all $0 \leq i \leq t$. Now, using the same arguments as above, we get $\psi^iH^t / \psi^{i+1}H^t = \varphi^iH^t / \varphi^{i+1}H^t$ for all $0 \leq i \leq t$ and so the result follows. $\square$

The following corollary extends [9, Corollary 3.3].

Corollary 2.3. *Let the situation be as in Theorem 2.2. Assume that N is Cohen-Macaulay with $\dim N = d$. Then $H_{\mathfrak{m}}^t(M, N) \cong \text{Ext}_R^t(M, N)$.*

Proof. By the same arguments as in the proof of Theorem 2.2 and [9, Corollary 3.3], we have $\text{Ext}_R^t(M, N) \cong \text{Ext}_R^{t-d}(M, H_{\mathfrak{m}}^d(N)) \cong E_{\infty}^{t-d,d}$,

$$0 = \phi^tH^t = \phi^{t-1}H^t = \dots = \phi^{t-d+1}H^t$$

and, also,

$$\phi^{t-d}H^t = \dots = \phi^0H^t = H_{\mathfrak{m}}^t(M, N)$$

for all t . It therefore follows that $H_{\mathfrak{m}}^t(M, N) \cong \text{Ext}_R^t(M, N)$. $\square$

The following theorem is related to [7, Theorem 1.2].

Theorem 2.4. *Let N be an R -module and t a positive integer. Then the following conditions are equivalent:*

- (i) $\text{Ext}_R^i(R/\mathfrak{a}, N)$ is Artinian for all $i \leq t$.
- (ii) $H_{\mathfrak{a}}^i(M, N)$ is Artinian for any finitely generated R -module M and all $i \leq t$.

Proof. (ii) $\implies$ (i) is immediate by [7, Theorem 1.2].

(i) $\implies$ (ii). By [7, Theorem 1.2] $H_{\mathfrak{a}}^i(N)$ is Artinian for all $i \leq t$. Now by similar arguments as in the proof of 1.1. One can see that $E_{\infty}^{i,t-i}$ and $\phi^i H^t$ are Artinian for all $0 \leq i \leq t$. We consider the exact sequences

$$0 \longrightarrow \phi^1 H^i \longrightarrow H_{\mathfrak{a}}^i(M, N) \longrightarrow E_{\infty}^{0,i} \longrightarrow 0 (0 \leq i \leq t).$$

Hence $H_{\mathfrak{a}}^i(M, N)$ is Artinian for all $i \leq t$. $\square$

Corollary 2.5. *Let the situation be as in Theorem 2.4. The following conditions are equivalent:*

- (i) $\text{Ext}_R^t(R/\mathfrak{a}, N)$ is Artinian for all t .
- (ii) $H_{\mathfrak{a}}^t(M, N)$ is Artinian for all finitely generated R -modules M and all t .

The following corollary is a generalization of [3, Theorem 2.2].

Corollary 2.6. *Let the situation be as in Corollary 2.5 and assume that $\dim(R/\mathfrak{a}) = 0$. Then $H_{\mathfrak{a}}^t(M, N)$ is Artinian for all finitely generated R -modules M and all t .*

REFERENCES

- [1] J. Asadollahi, K. Khashayarmanesh and Sh. Salarian, *On the finiteness properties of the generalized local cohomology modules*, Comm. Algebra., **30**(2002), 859-867.
- [2] M. H. Bijen-Zadeh, *A common generalization of local cohomology theories*, Glasgow Math. J. **21** (1980), 173-181.
- [3] K. Divaani-Aazar, R. Sazeeleh and M. Tousi, *On vanishing of generalized local cohomology modules*, Algebra Colloquium, to appear.
- [4] A. Grothendieck, *Local cohomology*, (notes by R. Hartshorne), LNM 41, springer, 1967.
- [5] J. Herzog, *Komplexe, Auflösungen und dualität in der lokalen Algebra*, Habilitationsschrift, Universität Regensburg. 1970.
- [6] C. Huneke, *Problems on local cohomology*, Free resolutions in commutative algebra and algebraic geometry, Res. Notes Math., **2** (1992), 93-108.

- [7] L. Melkersson, *A note on Artinian local cohomology modules*, priprint.
- [8] J. Rotman, *Introduction to homological algebra*, Academic Press, 1979.
- [9] P. Schenzel, *On the use of local cohomology in algebra and geometry*, Lectures at the summer school of commutative algebra and algebraic geometry, Ballaterra, 1996, Birkhäuser. Verlag, 1998.
- [10] N. Suzuki, *On the generalized local cohomology and its duality*, J. Math. Kyoto University, **18** (1978), 71-85.

A. MAFI, INSTITUTE OF MATHEMATICS, UNIVERSITY FOR TEACHER EDUCATION, 599
TALEGHANI AVENUE, TEHRAN 15614, IRAN AND DEPARTMENT OF MATHEMATIC, ARAK UNI-
VERSITY, ARAK, IRAN.

E-mail address: a-mafi@araku.ac.ir